Lamont– Doherty Earth Observatory
Office of Marine Affairs
61 Route 9W
Palisades, NY 10969

Prepared By: Richard Oliver–Goodwin
richardo@ldeo.columbia.edu
845 365–8677

R/V Maurice Ewing Data Reduction Summary

EW0306 Balboa, Panama − Bergen, Norway

Date	Julian Date	Time	Port
July 1, 2003	213	00:00:00	Balboa, Panama
August 19, 2003	231	08:39:40	Bergen, Norway



Project Summary

DESCRIPTION

Background and Scientific Objectives

Transit

Cruise Members

Science Party

None.

Ship's Science

John Byrne	Gunner	johnby@ldeo.columbia.edu
Ropate Maiwiriwiri	Core Bosun	ropate@ldeo.columbia.edu
Richard Oliver−Goodiwn	Computing	richardo@ldeo.columbia.edu
Justin Walsh	Gunner	cabinboy@ldeo.columbia.edu

Ship Crew

James O'Loughlin	Captain	captain@ewing.ldeo.columbia.edu
Albert Karlyn	Chief Engineer	engine@ewing.ldeo.columbia.edu
Dave Wolford	1st Mate	wolf@ewing.ldeo.columbia.edu
Rick Thomas	2nd Mate	rickt@ewing.ldeo.columbia.edu
Shankar Bhardwaj	3rd Mate	bhardwaj@ewing.ldeo.columbia.edu
Miguel Flores	1st A/Engineer	miguel@ewing.ldeo.columbia.edu
Matthew Ingerson	3rd A/Engineer	ingerson@ewing.ldeo.columbia.edu
Dexter James	3rd A/Engineer	dexter@ewing.ldeo.columbia.edu
Shawn Lindenmuth	Steward	shawn@ewing.ldeo.columbia.edu
David Philbrick	Bosun	davep@ewing.ldeo.columbia.edu
Sympongco, Arnold	A/B	arnold@ewing.ldeo.columbia.edu
Florendo, Rudy	Oiler	florendo@ewing.ldeo.columbia.edu
Moqo, Luke	Utility	luke@ewing.ldeo.columbia.edu
Von Mehren, Dennis	A/B	dennis@ewing.ldeo.columbia.edu
Lajoie, Richard	Oiler	lajoie@ewing.ldeo.columbia.edu
Marteny, Nicholas	O/S	marteny@ewing.ldeo.columbia.edu
Gahr, Edward	Cook	gahr@ewing.ldeo.columbia.edu
Uribe, Fernando	Oiler	gato@ewing.ldeo.columbia.edu
Strimback, Roger	O/S	

Cruise Notes

All data in this report is logged using GMT time and Julian days in order to avoid confusion with local time changes.

Hydrosweep

Hydrosweep acquisition was great during the transit with less than 2% average dropouts.

Gravity

No gravity notes

Magnetics

No data taken

Navigation

rt_nav was operated with gp04 as the primary GPS.

Time

no notes

Data Logging

The R/V Maurice Ewing data logging system is run on a Sparc Ultra Enterprise Server. Attached are 48 serial ports via 3 16−port Digi International SCSI Terminal Servers. Generally, all data logged by the Ewing Data Acquisition System (DAS) is time stamped with the CPU time of the server, and broadcast to the Ewing network using UDP packet broadcasts. The CPU time of the server is synchronized once every half hour to a Datum UTC gps time clock.

GPS times are also time−tagged with cpu time, although the time of the GPS position is from the GPS fix itself.

The following tables describe the data instruments which performed logging during this cruise. The tables associated with the instruments describe logging periods and data losses for that instrument.

Time Reference

Datum StarTime 9390−1000

logging interval: 30 minutes
file id: tr2

Used as the CPU synchronization clock. This clock is polled once every half hour to synchronize the CPU clock of the data logger to UTC time. The logger (octopus) is responsible for updating the times of the other CPUs.

This clock was running and synchronizing the system the entire cruise.

Interruption s greater than 30 minutes are displayed in the following table

Log Date	LogDate	Comment
2003+213:00:00:00.700		Logging officially started
2003+231:08:39:40.021		Logging officially ends

GPS Receivers

GPS data is usually logged at 1 − 2 second intervals. The NMEA strings GPGGA and GPVTG are logged for position, speed, and heading fixes. This data was logged constantly throughout the cruise.

Trimble Tasmon P/Y Code Receiver

logging interval: 2 seconds
file id: gp1

The Tasmon is no longer the primary GPS receiver for the Ewing Logging system. The accuracy is around 15 meters. There were no interruptions during this cruise.

Interruptions greater than 10 minutes are displayed in the following table

Log Date	LogDate	Comment
2003+213:00:00:00.410		Logging officially started
2003+231:08:39:39.278		Logging officially ends

Trimble NT300D

logging interval: 2 seconds
file id: gp2

The Trimble is the secondary receiver for GPS data. Data is logged at 2 second intervals and is also used as an input to Spectra, although it is weighed at a lower value than the Tasmon receiver.

Interruptions greater than 10 minutes are displayed in the following table

Log Date	LogDate	Comment
2003+213:00:00:00.112		Logging officially started
2003+216:15:39:03.342	2003+216:16:01:50.438	Data Interruption
2003+231:08:39:40.397		Logging Ends

C−Nav

logging interval: 2 seconds
file id: gp2

The C−Nav is a global satellite−based differential receiver. This is the best individual receiver currently on the ship.

Interruptions greater than 10 minutes are displayed in the following table

Log Date	LogDate	Comment
2003+213:00:00:02.531		Logging officially started
2003+231:08:39:39.479		Logging Ends

POS/MV

logging interval: 1 second
file id: gp4

The POS/MV is a receiver which uses C−Nav input, its own antennae, an inertial sensor, and optional RTCM corrections (when available) and a kalman filter to produce a smooth nav output and very accurate heading. As of June 2003 it is used as the primary GPS for Hydrosweep, as an input to Spectra, and can be used as the gps for reduction processing. With the C−Nav auxiliary input, this is the most accurate receiver on the ship.

Interruptions greater than 10 minutes are displayed in the following table

Log Date	LogDate	Comment
2003+213:00:00:00.000		Logging officially started
2003+231:08:39:40.021		Logging Ends

Speed and Heading

Furuno CI−30 Dual Axis Speed Log
Sperry MK−27 Gyro

logging interval: 6 seconds
file id: fu

The Furuno and Gyro are combined to output speed, heading and course information to a raw Furuno file, as well as an NMEA VDVHW signal used as an input to various systems including steering and Spectra.

Interruptions greater than 30 minutes are displayed in the following table

Log Date	Log Date	Comment
2003+213:00:00:01.629		Official start date
2003+231:08:39:39.058		Official end date

Gravity

Bell Aerospace BGM−3 Marine Gravity Meter System

logging interval: 1 second
file id: vc. (raw), vt. (processed)
drift per day: 0.175

The BGM consists of a forced feedback accelerometer mounted on a gyro stabilized platform. The gravity meter outputs raw counts approximately once per second which are logged and processed to provide real−time gravity displays during the course of the cruise as well as adjusted gravity data at the end of the cruise.

Interruptions greater than 10 minutes are displayed in the following table

Log Date	Log Date	Comment
2003+213:00:00:00.916		Official start date
2003+231:08:39:40.056		Official end time

Bathymetry

Krupp Atlas Hydrosweep−DS2

logging interval: variable based on water depth
file id: hb (centerbeam), hs (swath)

The hydrosweep full swath data is continuously logged for every cruise, and centerbeam data is extracted and processed separately. The centerbeam operates at a logging frequency dependent on the water depth.

The full swath data is not routinely processed, but can be processed with the MB−System software which can be downloaded for free. For instructions, use the website: http://www.ldeo.columbia.edu/MB−System.

MBSystem, version 5.0beta3 is necessary to process data after June 1, 2001.

Interruptions greater than 10 minutes are displayed in the following table

Log Date	LogDate	Comment
2003+213:00:00:00.000		Official start logging
2003+231:08:19:29.000		Official end logging

Weather Station

RM Young Precision Meteorological Instruments, 26700 series

logging interval: 1 minute
file id: wx

The weather station is used to log wind speed, direction, air temperature, and barometric pressure. We log this information at 1−minute intervals.

Log Date	LogDate	Comment
2003+213:00:00:00.001		Official start logging
2003+231:08:39:00.959		Official end logging

LOCATION 1

EW0305 Balboa, Panama

Pier/Ship	Latitude	Longitude
	08 57.011 N	079 34.017W

Pier 7, Balboa container teriminal

Reference	Latitude	Longitude
	8 57.325N	079 33.934W

Used tie from March 1995, between two nearby known reference points with low deviation between them.

	Id	Julian	Date	Mistie	Drift/Day	Prev Mistie
Pre Cruise	EW0304	186	05. Jul 03	24.64	−0.03	0.00
Post Cruise	EW0305	209	28. Jul 03	30.32	0.247	0.00
Total Days			23.00	5.68		

Time	Entry	Value	
17:35:00	CDeck Level BELOW Pier	2.50	
16:00:00	Pier 1 L&R Value	0.00	L&R
17:35:00	Reference L&R Value	0.00	L&R
17:45:00	Pier 2 L&R Value	0.00	L&R
	Reference Gravity	978223.50	mGals
	Gravity Meter Value (BGM Reading)	978256.30	mGals
	Needs Potsdam Correction	0	1 if Potsdam referenced

*Gravity meter is **5.5** meters below CDeck*

Difference in meters between Gravity Meter and Pier: 8.00 meters

Height Cor = Pier Height* FAA Constant

8.00	0.31			2.48 mGals/min

Difference in mGals between Pier and Gravity Meter

Pier (avg) − Reference *1.06 L&R/mGal — Delta L&R

0.00	0.00	1.06		0.00 mGals

Gravity in mGals at Pierside

Reference + Delta mGals [+ Potsdam] — IGSN−71 Referenced Pier (

978223.50	0.00	0.00		978223.50 mgals

Gravity in mGals at Meter

Pier Gravity+ Height Correction — Gravity@meter

978223.50	2.48			978225.98 mGals

Current Mistie

BGM Reading Calculated Gravity — Current Mistie

978256.30	978225.98			30.32 mGals

Location 2

EW0306 Bergen, Norway

Pier/Ship	Latitude	Longitude
	60 23.306N	005 18.556E

West end of Dokkeskjaerskaien (Container) Pier

Reference	Latitude	Longitude

No Lat/Lon available, (see map in gravity log).

	Id	Julian	Date	Mistie	Drift/Day	Prev Mistie
Pre Cruise	EW0305	209	28. Jul 03	30.32	0.25	5.68
Post Cruise	EW0306	240	28. Aug 03	35.74	0.175	0.00
Total Days			31.00	5.42		

Time	Entry	Value	
14:45:00	CDeck Level BELOW Pier	0.30	
14:45:00	Pier 1 L&R Value	5414.20	L&R
13:15:00	Reference L&R Value	5416.97	L&R
17:45:00	Pier 2 L&R Value	5414.20	L&R
	Reference Gravity	981951.10	mGals
	Gravity Meter Value (BGM Reading)	981972.10	mGals
	Needs Potsdam Correction	1	1 if Potsdam referenced

*Gravity meter is **5.5** meters below CDeck*

Difference in meters between Gravity Meter and Pier: 5.80 meters

Height Cor = Pier Height* FAA Constant

5.80	0.31			1.80 mGals/min

Difference in mGals between Pier and Gravity Meter

Pier (avg) − Reference *1.06 L&R/mGal = Delta L&R

5414.20	5416.97	1.06		−2.94 mGals

Gravity in mGals at Pierside

Reference + Delta mGals [+ Potsdam] = IGSN−71 Referenced Pier G

981951.10	−2.94	−13.60		981934.56 mgals

Gravity in mGals at Meter

Pier Gravity+ Height Correction = Gravity@meter

981934.56	1.80			981936.36 mGals

Current Mistie

BGM Reading − Calculated Gravity = Current Mistie

981972.10	981936.36			35.74 mGals

For all formats, a – in the time field means an invalid value for some reason.

Streamer Compass/Bird Data — cb.r

This data is not processed, but can still be found in the "processed" data directory.

```
Shot Time               Line     Shot     Latitude      Longitude

2000+079:00:08:40.085  strike1 000296  N 15 49.6217 W 060 19.8019

2nd GPS Position                      Tailbuoy Position
Latitude     Longitude                Latitude     Longitude

N 15 49.6189 W 060 19.8101    N 15 47.1234 W 060 20.1901

Furuno  Streamer
Gyro    Compasses & Heading

344.1       C01 2.3 C02 1.7  ...
```

Gun Depths — dg

Gun depths in tenths of meters. There will always be 20 gundepths even if only one gun was configured and shooting.

```
                        Gun Depths
Shot Time               1   2   3  4    5   6   7   8   9  ...  20

2001+089:06:47:05.909 189 068 005 005 096 005 060 054 005 ... 6
```

Raw Furuno Log — fu.s

This data has been smoothed and output 1 fix per minute.

```
CPU Time Stamp         Track Speed Hdg   Gyro

2000+166:00:01:53.091 –     4.4   140.5 148.3
```

Hydrosweep Centerbeam — hb.n

Hydrosweep data merged with navigation

```
                              Centerbeam
CPU Time Stamp       Latitude Longitude          Depth

2000+074:09:55:00.000  N 13 6.6206    W 59 39.3908   134.9
```

Merged Data — m

```
                                                        GPS
CPU Time Stamp         Latitude     Longitude          Used  Set  Drift Depth

2000+200:12:25:00.000  N 45 54.1583 W 42 47.1770       gp1   0.0  0.0

Magnetic                        Gravity
Total Intensity   Anomaly     FAA  GRV       EOTVOS   Drift   Shift

49464.7           55.5        22.2 980735.0   –8.4     –0.1     2.8

Temperature Salinity Conductivity

0.0         0.0      0.0
```

The gravity drift and shift are values that have been added to the raw gravity to make up for drift in the meter that has been lost in accordance with a gravity check at each port stop.

Temperature,Salinity and Conductivity will only be valid while logging a Thermosalinograph, which is not usually the case.

Magnetics Data — mg.n

- A minus sign in the time stamp is flagged as a spike point, probably noise...
- Anomaly is based on the International Geomagnetic Reference Field revision 2000

```
CPU Time Stamp        Latitude     Longitude      Raw Value Anomaly
200+077:00:23:00.000 N 16 11.2918 W  59 47.8258 36752.2   -166.8
```

Navigation File — n

```
CPU Time Stamp         Latitude     Longitude       Used  Set  Drift
2000+074:00:03:00.000 N 13 6.2214  W  59 37.9399   gp1   0.0   0.0
```

Navigation Block — nb0

Navigation is a compendium of Ewing logged data at shot time. The shot position here is the shot position from the Spectra system.

```
Shot Time               Shot # CPU Time               Shot Position
2001+088:00:00:00.606 016967 2001+088:00:00:03.031 N 30 11.8324 W 042 10.8162

Water  Sea    Wind   -------------Tailbuoy----------------- Line
Depth  Temp Spd  Dir Latitude     Longitude     Range Bearg Name   Speed Heading
2565.1 20.7 16.4 164 N 30 12.0427 W 042 14.7319 6296.3 93.5 MEG-10 4.2   101.1
```

Tailbuoy Navigation — tb1.c

Raw tailbuoy fixes

```
CPU Time Stamp         Latitude     Longitude       GPS Precision
2001+088:00:00:02.000 N 30 12.0424 W 042 14.7309   SA
```

GPS Precision is either SA, DIFF or PCODE

Ewing Processed Shot Times — ts.n

Shot times and positions based on the Ewing navigation data processing

```
CPU Time Stamp         Shot # Latitude     Longitude     Line Name
2000+079:00:08:01.507 000295 N 15 49.5703 W 060 19.7843 strike1
```

Shot Data Status — ts.n.status

The ts.nxxx.status file describes the line information for that day, giving some basic statistics about the line: start, end times; missing shots; start and end shots.

```
LINE   strike1: 98+079:00:00:15.568 : 000283 .. 002286
        MISSING: 347, 410, 1727
LINE   dip2: 98+079:23:05:22.899 : 000002 .. 000151
```

This example says that on Julian Day 079 of 1998, two lines (strike1 and dip2) were run: the end of strike 1 (shots 000283 to 002286) and the start of dip2 (shots 000002 to 000151).

Line strike1 had some missing shots in the data file (probably missing on the SEG-d header as well).

Spectra Shot Times — nb2.r

The shot times and positions based on the Spectra positioning; with raw tailbuoy range and bearing.

```
CPU Time Stamp          Shot # Latitude      Longitude      Line Name
2001+084:00:00:05.924 009245 N 23 31.2410 W 045 25.0894

                                 Tailbuoy
Latitude      Longitude      Range  Bearing  Line Name
N 23 30.4540 W 045 21.4338 6389.8 283.2    KANE-4
```

Raw Gravity Counts — vc.r

sample BGM-3 gravity count record (without time tag):

```
pp:dddddd ss
|    |    |___________ status: 00 = No DNV error; 01 = Platform DNV
|    |                         02 = Sensor DNV;   03 = Both DNV's
|    |________________ count typically 025000 or 250000
|_____________________ counting interval, 01 or 10
                       The input of data can be at 1 or 10 seconds.
```

Gravity Data — vt.n

* A minus sign in the time stamp is flagged as a spike point
* m_grv3 calculates the Eotvos correction as:
  eotvos_corr = 7.5038 * vel_east * cos(lat) + .004154 * vel*vel
* The theoretical gravity value is based upon different models for the earth's shape.
  - 1930 = 1930 International Gravity Formula
  - 1967 = 1967 Geodetic Reference System Formula
  - 1980 = 1980 Gravity Formula
* The FAA is computed as:
  faa = corrected_grv – theoretical_grv
* Velocity smoothing is performed w/ a 5 point window

```
CPU Time Stamp          Latitude      Longitude      Model FAA      RAW
2000+148:00:10:00.000 N 09 34.7255 W 085 38.5826 1980  9.48    978264.16

Eotvos  Drift DC      Raw  Velocity     Smooth Velocity
Smooth  Total Shift   North     East     North    East
-74.78  0.06  4.16    1.875 -10.373     1.927 \10.166
```

Datum Time — ts2.r

```
CPU Time                Datum Time        Time Reference
2001+069:00:15:29.727 069 00 15 29.378 datum
```

Raw GPS — gp(12).d, tb1.d

Raw GPS is in NMEA Format.

Meteorological Data wx

```
                         True
CPU Time Stamp          Spd Dir
2001+045:00:00:00.967   7.8  22

Bird1:                                       Bird 2
Speed                   Direction            Speed                  Direction
Inst  60sA   60mA  60sM  Inst 60sA 60mA      Inst  60sA  60mA  60sM   Inst 60sA 60mA

7.8   6.6    8.5   16.8  277  291  5         0.0   0.0   0.0   0.0    0    0    0

Temperature                    Humidity
Inst  60mA  60mm  60mM         Inst 60mm 60mM        Barometer

15.0  14.2  14.3  15.1         92   90   93          1027.5

Inst:       Current
60sA:       60 second average
60mA:       60 minute average
60sM:       60 second maximum
60mm:       60 minute minimum
60mM:       60 minute maximum
```

Merged Meteorological Data mmet

```
TSG, WX, CT merged with Nav at 1 minute fixes
date       time       lat      lon      gpu head spd
2001+244:00:00:00.000 12.14071 44.98469 gp1 10.2 83.0

tws  twd   temp hum  press  cti  cte  con sal  ct
26.5 228.0 30.6 87.0 1000.8 28.8 28.8 5.9 36.3 28.8

gpu  = gps unit in use
head = ship's heading
spd  = ship's speed in knots
tws  = true wind speed
twd  = true wind direction
temp = air temp (celcius)
hum  = relative humidity (%)
press= pressure in mb
cti  = sea temp from the internal TSG sensor
cte  = sea temp from the external TSG sensor
con  = conductivity, Siemens/meter
sal  = salinity, practical salinity units
ct   = sea temp from the C-keel sensor (to tenths of a degree)
```

Tape Contents

EW0306/	
EW0306.pdf	this document
ew0306.cdf	NetCDF database file of this cruise
ew0306.cdf_nav	NetCDF database file of this cruise' navigation
docs/	File Formats, Spectra manuals
processed/	Processed datafiles merged with navigation
trackplots/	daily cruise track plots (*postscript*)
raw/	Raw data directly from logger
reduction/	Reduced data files
clean/	daily processing directory, includes daily postscript plots of the data.