

---

# **LMG0407 - Shuttle**

## **Cruise HazWaste Run**

By W. Kevin Pedigo

**June 12, 2004 – June 27, 2004**

# Table of Contents

<b>INTRODUCTION .....</b>	<b>1</b>
<b>ARCHIVE DATA EXTRACTION .....</b>	<b>2</b>
<b>CD DIRECTORY STRUCTURE.....</b>	<b>3</b>
<b>DISTRIBUTION CONTENTS.....</b>	<b>4</b>
ADCP .....	4
CALIBRATION.....	4
CTD .....	4
/CTD/raw.....	4
/CTD/process.....	4
CRUISE TRACK .....	5
ICE IMAGES .....	5
ISOBAR CHARTS .....	5
DATA AND SCIENCE REPORT .....	5
QC PLOTS .....	5
XBT .....	5
/XBT/.....	5
<b>SCIENCE AND MOORING DATA.....</b>	<b>5</b>
JGOFS DATA SET.....	6
RVDAS .....	6
<i>Meteorological and Light Data</i> .....	7
<i>Navigational Data</i> .....	7
<i>Geophysical Data</i> .....	7
<i>Oceanographic Data</i> .....	7
<b>DATA FILE NAMES AND STRUCTURES.....</b>	<b>8</b>
LKNU.....	8
LMET.....	8
LTSG.....	9
LPCO.....	9
SVP1.....	9
LADC.....	9
LASH.....	10
LGYR.....	10
TGPS.....	11
LPCD.....	13

---

<b>LMG SENSORS .....</b>	<b>14</b>
SHIPBOARD SENSORS .....	16
CTD SENSORS .....	<b>ERROR! BOOKMARK NOT DEFINED.</b>
<b>ACQUISITION AND PROCESSING INFORMATION .....</b>	<b>17</b>
PROCESSING SPECIFICS .....	17
ERRORS AND EVENTS .....	17

## Introduction

The LMG data acquisition systems continuously log data from a suite of instrumentation throughout the cruise. This document describes the format of that data and its location on the distribution CDs. It also contains important information which may affect how this data is processed such as instrument failures or other known problems with acquisition.

The data collected during this cruise is distributed on a CD-ROM written in ISO9660 level-1 format. This data format has very strict requirements on filenames and organization. However, it is readable by virtually every computing platform.

All of the data has been archived with the Unix "tar" command and/or compressed using Unix "gzip" compression. Tar files have a ".tar" extension and Gzipped files have a ".gz" extension. Tools are available on all platforms for uncompressing and de-archiving these formats. On Macintosh, Stuffit Expander with DropStuff will open a tar archive and uncompress gzipped and Unix compressed files. For Windows9X, WinZip, a shareware utility included on this CD (remember, it is shareware) will open these files.

In some cases to adhere to the ISO9660 format the .tar extension was removed. When we tarred the files then gzip the tar archive the name of the file became *File.tar.gz*. This name does not follow the 8.3 naming convention of the ISO9660 format. So the file was renamed without .tar making the file name **File**.gz. On Windows and Mac Platforms Winzip and Stuffit Expander handles this just fine. When they expand the *File.gz* the expanded file becomes *File.tar*, which both software packages can handle. On Unix platform gunzip expands *File.gz* but it does not append the .tar extension. So you may not recognize the file as a tar archive, but OS does recognize it as a tar archive. If you use the file command it will return saying it is a tar file. The below tar command will un-archive the file just fine.

***IMPORTANT: Read the last section in this document, Acquisition Problems and Events, for important information that may affect the processing of this data.***

## Archive Data Extraction

It is often useful to know exactly how an archive was produced when expanding its contents. Tar files were created using the following commands:

```
tar cvf archive-file files-to-be-archived
```

To create a list of the files in the archive:

```
tar tvf archive-file > contents.list
```

To extract the files from the archive:

```
tar xvf archive-file file(s)-to-extract
```

G-zipped files will have a “.gz” extension on the filename. These files can be decompressed after de-archiving, using:

```
gunzip filename.gz
```

## CD Directory Structure

**/ADCP/**

Pingdata/

Gentoo/

**/CAL/**

SVP\_CALS.tar

UW\_CALS.tar

CTD\_CALS.zip

MET\_CALS.zip

Instrument.coeff

**/CTD/**

Raw/

Process/

**/ICE\_IMAGE/**

\*.jpg

**/ISOBARS/**

\*.gif

**/JGOF/**

Imgjgof.tar

**/MAPS/**

\*.jpg

\*.ps

Track.txt

**/MOORING/**

Moor2003.zip

**/QC\_PLOTS/**

Lmgqc.tar

**/REPORT/**

Report.doc

Report.html

Report.txt

**/RVDAS/**

Lmguw.tar

Lmgnav.tar

**/SCIENCE/**

Report/

Met\_Data/

**/SITREPS/**

\*.txt

**/UTILITY/**

Winzip

Stuffit Expander

**/XBT/**

\*.EDF

\*.RDF

## Distribution Contents

### ADCP

/Adcp/

The ADCP DAS data files are named pingdata.xxx (xxx representing a file number). Note that these extensions do NOT represent Julian day numbers. Please refer to the file's creation date. The ADCP DAS computer creates a new pingdata file when the current one reaches a size of 320K. The ping files logged on Gentoo, the Linux processing computer, however are created new each day.

Some ADCP data is also transmitted to RVDAS. East and North vectors for ship's speed relative to the reference layer and ship's heading are archived in the navigational data section of RVDAS.

### Calibration

/Cal/

The tar files in the Cal directory contain images of calibration sheets for each of the following systems: Sound Velocity Probe(SVP\_CALS.TAR), Meteorological System(MET\_CALS.TAR), Underway System(UW\_CALS.TAR), and CTD(CTD\_CALS.TAR).

CTD diagrams can be found in the ctd Cal directory for voltage and frequency sensors.

### CTD

/CTD/

The ctd data was collected and processed on a windows 98 computer, using Seasave Win32 – Version 5.25a and SBE Data Processing – Version 5.21.

For more information and software visit the web site at [www.seabird.com](http://www.seabird.com).

/CTD/raw

This directory contains the original .con, .dat, .bl, and .hdr files files produced for each CTD cast.

/CTD/process

Each processed cast is represented here by a set of ten files. Note that cast 1 was split into a separate data set for the up and down segments. Also, the DO2 sensor was only added onto to the instrument package for the 4<sup>th</sup> cast. In addition to the standard files listed below, an additional round of post processing was done to produce rg302EEE.cnv and .asc files containing non-bin averaged data and an ascii conversion of that data.

G404EEE.con	A copy of the configuration file for the cast.
G404EEE.cnv	The converted file for the whole cast.
G404EEE.ros	The rosette file that contains the scan lines for each bottle trip.
G404EEE.btl	The bottle file that contains the avg, standard deviation, min, and max for a select set of variables for each bottled fired during the upcast.
Dg404EEE.cnv	The converted file for the down cast.
Dg404EEE.asc	An ASCII formatted file for the down cast without a header.
Dg404EEE.hdr	The header for the down cast.
Ug404EEE.cnv	The converted file for the up cast.
Ug404EEE.asc	An ASCII formatted file for the up cast without a header.
Ug404EEE.hdr	The header for the up cast.

## **Cruise Track**

/MAP/

PostScript cruise track plots have been produced for this cruise, in three sizes. It has also been converted to a jpeg format. The cruise track file track.txt is also included and contains the latitude and longitude in 10-minute intervals.

## **Ice Images**

/ICE\_IMAGE/

This directory contains .jpg files of Terrascan ice imagery sent to the ship from Palmer station before and during the cruise.

## **Isobar Charts**

/Isobars/

This directory contains GIF image files. These file are an analysis of mean sea level pressure from the National Center for Environmental Prediction's Medium Range Forecast Model. They are updated every 6 hours. Naming the convention is as follows yyjjj.hh.gif where yy is the year, jjj is the day number, and hh is the hour.

## **Data and Science Report**

/Report/

Copies of this report in MS Word, HTML, and text formats.

## **QC Plots**

/QC\_PLOTS/

Postscript files of data stored each day on RVDAS for quality control analysis during the cruise. There are 3 types of files, named metXXX.ps, navXXX.ps, and oceanXXX.ps, where XXX is represents the Julian day. Met files are a summary of the data from the meteorological instruments, Nav files are a summary of navigational data, and Ocean files are a summary of the underway seawater and bathymetry data.

## **XBT**

/XBT/

Expendable Bathythermographic (XBT) probes were used to obtain water column temperature profiles. These XBT were launched at all SIO Mooring sites. The .RDF files contain the raw data, and the .EDF contained the exported ascii data.

## **Science and Mooring Data**

/Science/ and /Moor2003/



These directories contain additional data provided by the PI to supplement the standard distribution data set.

## JGOFS Data Set

/JGOF/

The JGOFS data set consists of a single file produced each day named jg<julian\_day>.dat.gz where <julian\_day> is the day the data was acquired. The “.gz” extension indicates that the individual files are compressed before archiving. The daily file consists of 22 separate columnar fields in text format, which are described below. The JGOFS data set is obtained primarily by applying calibrations to raw data and decimating to whole minute intervals. However, several fields are derived measurements from more than a single raw input. *Note: Null, unused, or unknown fields are filled with 9's in the JGOFS data.*

Additionally, 3 separate QC plots are generated daily by the ET using the JGOFS data set. These plots include TSG and Bathymetry data, meteorological data, and navigation data. The files are called ocean<julian\_day>.ps, met<julian\_day>.ps, and nav<julian\_day>.ps respectively.

Field	Data	Units
01	GMT date	dd/mm/yy
02	GMT time	hh:mm:ss
03	PCOD latitude (negative is South)	Ddd.dddd
04	PCOD longitude (negative is West)	Ddd.dddd
05	Ships speed	Knots
06	GPS HDOP	-
07	Gyro Heading	Degrees (azimuth)
08	Course over ground	Degrees (azimuth)
09	Mast PAR	μEinsteins/meters <sup>2</sup> sec
10	Sea surface temperature	°C
11	Not used	-
12	Sea surface salinity	PSU
13	Sea depth (uncorrected, calc. sw sound vel. 1500 m/s)	meters
14	True wind speed (port windbird)	meters/sec
15	True wind direction (port windbird)	degrees (azimuth)
16	Ambient air temperature	°C
17	Relative humidity	%
18	Barometric pressure	mBars
19	Sea surface fluorometry	volts (0-5 FSO)
20	Not used	-
21	PSP	W/m <sup>2</sup>
22	PIR	W/m <sup>2</sup>

## RVDAS

/rvdas/

RVDAS (Research Vessel Data Acquisition System) was developed at Lamont-Doherty Earth Observatory of Columbia University and has been used on the R/V Maurice Ewing for several years. It was adapted for use on the Nathaniel B. Palmer and her sister ship, the R/V Laurence M. Gould.

Below you will find detailed information on the data included. Be sure to read the “Significant Acquisition Events” section below for important information about data acquisition during this cruise.

*Meteorological and Light Data*

Measurement	File ID	Collect. Status	Rate	Instrument
Air Temperature	lmet	continuous	1 sec	R. M. young 41372VC
Relative Humidity	lmet	continuous	1 sec	R. M. young 41372VC
Wind Speed/Direction	lmet	continuous	1 sec	R. M. young 5106
PIR (LW radiation)	lmet	continuous	1 sec	Eppley PIR
PSP (SW radiation)	lmet	continuous	1 sec	Eppley PSP
Photosynthetically-Available Radiation	lmet	continuous	1 sec	BSI QSR-240
Barometer	lmet	continuous	1 sec	R. M. young 61201

*Navigational Data*

Measurement	File ID	Collect. Status	Rate	Instrument
Attitude GPS	lash	continuous	1 sec	Ashtec ADU-2
P-Code GPS	lpcd	Not collected	1 sec	Trimble 20636-00SM
Gyro	lgyr	continuous	0.2 sec	Anschutz Gyro
Trimble GPS	tgps	continuous	1 sec	NT200

*Geophysical Data*

Measurement	File ID	Collect. Status	Rate	Instrument
Bathymetry	lknu	variable	Varies	Knudsen 320B/R

*Oceanographic Data*

Measurement	File ID	Collect. Status	Rate	Instrument
Salinity	ltsg	continuous	15 sec	SeaBird 21
Sea S Temperature	ltsg	continuous	15 sec	SeaBird 3-01/S
Fluorometry (analog)	ltsg	continuous	15 sec	Turner 10-AU-005
ADCP	ladc	continuous	1 sec	RD Instruments

## Data File Names and Structures

RVDAS data is divided into two broad categories, **Underway** and **Navigation**. The groups are abbreviated “uw” and “nav”. Thus, these two tar files, lmguw.tar and lmgnav.tar exist under the top-level rvdas directory. The instruments are broken down as shown. Each data file is g-zipped to save space on the distribution. Not all data types are collected everyday or on every cruise.

RVDAS data files are named following the convention: LMG[FileID].dDDD.

- The FileID is a 4-character code representing the system being logged, for example: lmet (for meteorology)
- DDD is the Julian day of the data collection

Underway Data	File ID	Navigation Data	File ID
Meteorological	lmet	Gyro Compass	lgyr
Knudsen	lknu	P-CODE GPS	lpcd
Thermosalinograph	ltsg	Ashtech ADU2 GPS	lash
ADCP	ladc	Trimble NT2100 GPS	tgps
Sound Velocity Probe	lsvp	PCO2 System	lpcd

Data is received by the RVDAS system via RS-232 serial connections. The data files that comprise the rvdas data set are described below. A time tag is added to each line of data received and the data is written to disk.

```
YY+DDD:HH:MM:SS.SSS [data stream from instrument]
```

Where, YY: two-digit year, DDD: Julian Day, HH: 2 digit hours, MM: 2 digit minutes SS.SSS: seconds. All times are UTC.

The delimiters used to separate fields in the raw data files are usually spaces and commas, but other delimiters are used (:, =, @) and occasionally there is no delimiter. Care should be taken when reprocessing the data that the fields separations are clearly understood. An example data

### lknu

```
99+099:00:18:19.775 hf,305.2,lf,304.3
```

Field	Data	Units
1	RVDAS Time Tag	
2	hf – high frequency flag (12 kHz)	
3	high frequency depth	meters
4	lf – low frequency flag (3.5 kHz)	
5	low frequency depth	meters

### lmet

```
02+314:23:59:50.067 01.2 047 028 01.3 063 042 0988.8 001.7 084 -000.2192 0000.9358 0025.5875
```

Field	Data	Units
1	RVDAS Time Tag	
2	Port Wind Speed	m/s
3	Port Wind Direction	deg
4	Port Wind Direction (standard deviation)	deg
5	Starboard Wind Speed	m/s
6	Starboard Wind Direction	deg
7	Starboard Wind Direction (standard deviation)	deg
8	Barometer	millibars
9	Temperature	°C
10	Relative humidity	%
11	PSP (long wave radiometer)	Volts

Field	Data	Units
12	PIR (short wave radiometer)	Volts
13	PAR (photo-synthetically available radiation, 400 - 700 nm)	Volts

**Itsg**

02+310:23:57:30.200 8542 -1.2580 34.1740 -1.2030 3.435 0.349 27.361205

Field	Data	Units
1	RVDAS Time Tag	
2	Scan number	
3	Internal water temperature	°C
4	Salinity	PSU
5	External water temperature	°C
6	Transmissometer signal	Volts
7	Fluorometer signal (analog)	Volts
8	Conductivity	S/cm

**Ipco**02+319:23:59:13.748 2002319.99851 7154.27 26.49 1033.6 325.79 6.74 329.3  
53.76 0 Equil

Field	Data	Units
1	RVDAS Time Tag	
2	Julian date file string	Julian
3	IR voltage reading	mV
4	Cell temperature	°C
5	Barometer	millibars
6	VCO2	mL
7	Equilibrator temperature	°C
8	PCO2	millibars
9	Gas flow	mL/min
10	Solenoid position ID	number
11	Valve Position ID	number
12	Measured gas	name

**svp1**

00+348:01:59:52.128 1539.40

Field	Data	Units
1	RVDAS Time Tag	
2	Sound velocity	m/s

**ladc**

00+019:23:59:59.099 \$PUHAW,UVH,-1.48,-0.51,250.6

Field	Data	Units
1	RVDAS Time Tag	
2	\$PUHAW	
3	UVH (E-W, N-S, Heading)	
4	Ship Speed relative to reference layer <sup>1</sup> velocity <sup>2</sup> , East vector	knots
5	Ship Speed relative to reference layer <sup>1</sup> velocity <sup>2</sup> , North vector	knots
6	Ship heading	degrees

<sup>1</sup>The reference layer is an average velocity measured in a number of depth “bins”. On the LMG, the bins are eight meters deep and bins 3-10 define the reference layer. Hence, the reference layer is the water column from 16-80 meters beneath the ship.

<sup>2</sup>The speed output is water velocity relative to the ship’s hull and is therefore opposite of the actual movement of the ship. For example, if the ship’s heading is due north, the North/South reference layer velocity is likely to be negative (southerly).

## lash

### ATTD: Attitude Data

01+081:00:00:00.806 \$PASHR,ATT,345605.0,165.03,+001.86,-01.96,0.0018,0.0173,0\*22

Field	Data	Units
1	RVDAS Time Tag \$PASHR	
2	ATT	
3	GPS Time sec. of the week	seconds
4	heading (rel. to true North)	degrees
5	pitch	degrees
6	roll	degrees
7	Measurement RMS error	meters
8	Baseline RMS error	meters
9	attitude reset flag	

01+081:00:00:00.966 \$GPGGA,235952.00,6051.7937,S,06030.2175,W,1,08,01.0,+00068,M,,M,,\*79

Field	Data	Units
1	RVDAS Time Tag \$GPGGA	
2	UTC time at position	hhmmss.ss
3	Latitude	ddmm.mmm
4	North (N) or South (S)	
5	Longitude	ddmm.mmm
6	East (E) or West (W)	
7	GPS quality (1=GPS 2=DGPS)	
8	Number of GPS satellites used	
9	HDOP	
10	Antenna Height	meters
11	M for Meters	
12	Geoidal height	meters
13	M for meters	
14	age of diff. GPS data	sss
15	differential reference station ID	aaaa

## lgyr

02+315:23:59:58.194 \$PASVW,00.1,A\*1D

02+315:23:59:58.414 \$IIVHW,287.7,T,,M,,N,,K\*71

02+315:23:59:58.616 \$HEHDT,287.7,T\*25

02+315:23:59:58.821 \$HEROT,001.6,A\*2C

02+315:23:59:58.984 \$HCHDT,,T\*07

### HDT: True Heading

01+083:00:00:02.893 \$HEHDT,246.3,T\*2C

Field	Data	Units
-------	------	-------

Field	Data	Units
1	RVDAS Time Tag \$HEHDT	
2	Heading XXXXX = ddd.d	degrees
3	T flag for true heading, checksum	

## ROT: Rate of Turn

01+083:00:00:03.093 \$HEROT,-006.3,A\*03

Field	Data	Units
1	RVDAS Time Tag \$HEROT	
2	Rate of turn	degrees/min
3	Status: A = data valid, checksum	

## tgps

## GGA: Global Positioning Fix Data

00+040:00:00:00.985 \$GPGGA,000003,6139.961,S,05949.422,W,1,6,001.64,-00036,M,00000,M,,

Field	Data	Units
1	RVDAS Time Tag \$GPGGA	
2	Latitude in degrees with decimal minutes	ddmm.mmm
3	North (N) or South (S)	
4	Longitude in degrees with decimal minutes	ddmm.mmm
5	East (E) or West (W)	
6	GPS quality (1=GPS 2=DGPS)	
7	Number of GPS satellites used	
8	Horizontal dilution of precision (HDOP)	
9	Antenna height above/below mean-sea-level (geoid)	meters
10	Units for antenna height (M = Meters)	
11	Geoidal Separation <sup>1</sup>	
12	Units for Geoidal Separation (M = Meters)	meters
13	Age of differential GPS data, number of seconds since last SC104 Type 1 or 9	
14	Differential reference station ID	

<sup>1</sup>Geoidal Separation: the difference between the WGS-84 earth ellipsoid and mean-sea-level (geoid). A negative value represents mean-sea-level below ellipsoid.

## GLL: Geographic Position – Latitude/Logitude

00+040:00:00:00.065 \$GPGLL,6139.96,S,05949.42,W,000002,A

Field	Data	Units
1	RVDAS Time Tag \$GPGLL	
2	Latitude	ddmm.mmm
3	North (N) or South (S)	
4	Logitude	ddmm.mmm
5	East (E) or West (W)	
6	UTC of position	hhmmss.ss
7	Status: A = Data Valid	

## VTG: Track Made Good and Speed over Ground

00+040:00:00:00.213 \$GPVTG,161,T,149,M,009.6,N,017.8,K

Field	Data	Units
1	RVDAS Time Tag \$GPVTG	
2	Track, degrees true	degrees
3	T flag for True	
4	Track, degrees magnetic	degrees
5	M flag for Magnetic	
6	Speed over Ground	knots
7	N flag for Knots	
8	Speed over Ground	km/hr
9	K flag for km/hr	

## VHW: Speed Through Water and Heading

00+040:00:00:00.212 \$GPVHW,246,T,234,M,012.3,N,022.8,K

Field	Data	Units
1	RVDAS Time Tag \$GPVHW	
2	Heading, degrees True	degrees
3	T flag for True	
4	Heading, degrees Magnetic	degrees
5	M flag for Magnetic	
6	Speed through water	knots
7	N flag for Knots	
8	Speed through water	km/hr
9	K flag for km/hr	

## ZDA: Time and Date

00+040:00:00:00.285 \$GPZDA,000002,09,02,2000,00,00

Field	Data	Units
1	RVDAS Time Tag \$GPZDA	
2	UTC time	hhmmss.ss
3	Day: 01 – 31	dd
4	Month: 01 – 12	mm
5	Year	yyy
6	Local time zone description <sup>1</sup> , 00 +/-13 hrs	
7	Local time zone minutes description, same sign as local hours	

<sup>1</sup>Zone description is the number of whole hours added to local time to obtain GMT, values are negative for East longitudes.

### BWC: Bearing and Distance to Waypoint

00+040:00:00:00.865 \$GPBWC,000003,6209.70,S,05824.00,W,127.2,T,115.3,M,050.1,N,014

Field	Data	Units
1	RVDAS Time Tag \$GPBWC	
2	UTC of bearing	hhmmss.ss
3	Destination waypoint latitude in degrees, decimal minutes	ddmm.mmm
4	Hemisphere Flag: N or S	
5	Destination waypoint longitude in degrees, decimal minutes	ddmm.mmm
6	Hemisphere Flag: E or W	
7	Bearing, degrees true	degrees
8	T flag for True	
9	Bearing, degrees magnetic	degrees
10	M flag for Magnetic	
11	Distance to waypoint in nautical miles	nm
12	N flag for Nautical Miles	
13	Waypoint ID	

### Ipcd

#### GGA: GPS Position Fix – Geoid/Ellipsoid

00+019:23:59:59.301 \$GPGGA,235958.409,6849.6944,S,13712.8472,W,1,06,1.2,092.4,M,047.3,M,,\*67

Field	Data	Units
1	RVDAS Time Tag \$GPGGA	
2	UTC time at position	hhmmss.sss
3	Latitude	ddmm.mmm
4	North (N) or South (S)	
5	Longitude	ddmm.mmm
6	East (E) or West (W)	
7	GPS quality (1=GPS 2=DGPS 3=P-CODE)	
8	Number of GPS satellites used	
9	HDOP	
10	Antenna Height	meters
11	M for Meters	
12	Geoidal height	meters
13	M for meters	
14	Null field	
15	Checksum	



## GLL: GPS Latitude/Longitude

00+019:23:59:59.381 \$GPGLL,6849.6944,S,13712.8472,W,235958.409,A\*35

Field	Data	Units
1	RVDAS Time Tag \$GPGLL	
2	Latitude	degrees
3	North or South	
4	Longitude	degrees
5	East or West	
6	UTC of position	hhmmss.sss
7	status of data (A = valid)	
8	Checksum	

## VTG: GPS Track and Ground Speed

00+019:23:59:59.382 \$GPVTG,238.7,T,182.3,M,001.8,N,003.3,K\*41

Field	Data	Units
1	RVDAS Time Tag \$GPVTG	
2	Heading	degrees
3	degrees True (T)	
4	Heading	degrees
5	degrees magnetic (M)	
6	Ship speed	knots
7	N = knots	
8	Speed	km/hr
9	Checksum	

## Ifir

## Digital output of Turner Fluorometer

04+107:16:48:02.342 0 5450 :: 4/16/04 09:44:17 = 0.632 (RAW)  
 04+107:16:48:03.342 0 5451 :: 4/16/04 09:44:18 = 0.631 (RAW)  
 04+107:16:48:04.382 0 5452 :: 4/16/04 09:44:19 = 0.631 (RAW)  
 04+107:16:48:05.382 0 5453 :: 4/16/04 09:44:20 = 0.631 (RAW)

Field	Data	Units
1	RVDAS Time Tag	
2	Zero Field	numeric
3	Sample Number	numeric
4	Fluorometer Date	mm/dd/yy
5	Fluorometer Time	hh:mm:ss
6	Digital output of fluorometer	Volts
7	(RAW)	

**loxyg**

PCO2 system Oxygen sensor, a TD218 Oxygen Optode 330.

For further information on this data, contact Tim Newberger at [tnewberg@ldeo.columbia.edu](mailto:tnewberg@ldeo.columbia.edu)

```

04+117:23:57:23.504 MEASUREMENT      3830      380 Oxygen:      309.95      Saturation:
83.48 Temperature:      -1.35      DPhase:      33.41      BPhase:      32.22
      RPhase:      0.00      BAmp:      262.09      BPot:      163.00      RAmp:
0.00 RawTem.:      694.92
04+117:23:58:23.508 MEASUREMENT      3830      380 Oxygen:      309.59      Saturation:
83.38 Temperature:      -1.35      DPhase:      33.43      BPhase:      32.23
      RPhase:      0.00      BAmp:      262.14      BPot:      163.00      RAmp:
0.00 RawTem.:      694.95
04+117:23:59:23.512 MEASUREMENT      3830      380 Oxygen:      309.74      Saturation:
83.43 Temperature:      -1.35      DPhase:      33.42      BPhase:      32.22
      RPhase:      0.00      BAmp:      262.07      BPot:      163.00      RAmp:
0.00 RawTem.:      694.83

```

Field	Data	Units
1	RVDAS Time Tag	
2-4	Measurement ID, Model Number, Serial Number	alphanumeric
5	Oxygen heading	text
6	Oxygen Reading	Raw numeric
7	Saturation heading	text
8	Saturation Reading	Raw numeric
9	Temperature heading	text
10	Water Temperature	°C
11	Dphase heading	text
12	Dphase	Raw numeric
13	Bphase heading	text
14	BPhase	Raw numeric
15	Rphase heading	text
16	Rphase	Raw numeric
17	Bamp heading	text
18	Bamp	Raw numeric
19	Bpot heading	text
20	Bpot	Raw numeric
21	Ramp heading	text
22	Ramp	Raw numeric
23	RawTem heading	text
24	RawTemp	Raw numeric

## LMG Sensors

### Shipboard Sensors

Sensor	Description	Serial #	Cal. Date	Status
Port Anemometer	R.M. Young 105106	WM35061	25-Jun-03	collected
Stbd Anemometer	R.M. Young 105106	WM5708	25-Jun-03	collected
Barometer	R.M. Young 61201	BP01150	26-Feb-03	collected
Humidity/Wet Temp	R.M. Young 41372VC	41372LC	23-Dec-03	collected
Mast PAR	BSI QSR-240P	6393	12-Dec-03	collected
Pyranometer	Eppey PSP	28933F3	30-Mar-04	collected
Pyrgeometer	Eppey PIR	32031F3	21-Nov-03	collected
TSG	SeaBird SBE21	1789	12-Sep-03	collected
TSG Remote Temp	SeaBird 3-01/S	031619	21-Oct-03	collected
Fluorometer	Turner 10-AU-005 Lamp: daylight 10-045, reference filter: 10-052, emission filter: 10-051, excitation filter: 10-050.	6592RTX	n/a	collected
Transmissometer	WET Labs C-Star 25cm	CST424PR	06-Nov-03	collected
P-Code GPS	Trimble 20636-00 (SM)	220035265	n/a	collected
Bathymetry	Knudsen 320B/R	K97715	n/a	collected

## Acquisition and Processing Information

### Processing Specifics

Refer to the instrmnt.cof file along with the specific instrument calibration sheets, both located in the Cal/ directory of the data distribution, for information on how the RVDAS data was collected and processed.

### Errors and Events

This section lists all significant events and known problems with acquisition during this cruise including instrument failures, data acquisition system failures, and other factors affecting this data set.

Date (Julian)	Time (GMT)	Event	Location
165	00:26	START DAS and all instruments	68° West
165	19:02	Turned off ADCP bottom tracking	55°01'S, 64°56'W
167	20:26	Turned on ADCP bottom tracking	62°29'S, 62°25'W
168	16:36	Disabled Acoustic/Seawater Sys	Arrive Palmer Station.
175	14:00	Enabled Acoustic/Seawater Sys	Depart Palmer Station.
178	21:21	STOP DAS and all instruments	68° West